

NGC 3576 and NGC 3603: Two Luminous Southern HII Regions Observed at High Resolution with the Australia Telescope Compact Array

C. G. De Pree¹, Melissa C. Nysewander^{1,2}, W. M. Goss²

¹Department of Physics & Astronomy, Agnes Scott College, 141 E. College Ave, Decatur, GA 30030

²National Radio Astronomy Observatory, P.O.Box 0, Socorro, NM 87801

ABSTRACT

NGC 3576 (G291.28-0.71; $l = 291.3^\circ$, $b = -0.7^\circ$) and NGC 3603 (G291.58-0.43; $l = 291.6^\circ$, $b = -0.5^\circ$) are optically visible, luminous HII regions located at distances of 3.0 kpc and 6.1 kpc, respectively. We present 3.4 cm Australian Telescope Compact Array (ATCA) observations of these two sources in the continuum and the H90 α , He90 α , C90 α and H113 β recombination lines with an angular resolution of $7''$ and a velocity resolution of 2.6 km s^{-1} . All four recombination lines are detected in the integrated profiles of the two sources. Broad radio recombination lines are detected in both NGC 3576 ($\Delta V_{FWHM} \geq 50 \text{ km s}^{-1}$) and NGC 3603 ($\Delta V_{FWHM} \geq 70 \text{ km s}^{-1}$). In NGC 3576 a prominent N-S velocity gradient ($\sim 30 \text{ km s}^{-1} \text{ pc}^{-1}$) is observed, and a clear temperature gradient (6000 K to 8000 K) is found from east to west, consistent with a known IR color gradient in the source. In NGC 3603, the H90 α , He90 α and the H113 β lines are detected from 13 individual sources. The Y^+ (He/H) ratios in the two sources range from 0.08 ± 0.04 to 0.26 ± 0.10 . The H113 β /H90 α ratio in NGC 3576 is close to the theoretical value, suggesting that local thermodynamic equilibrium (LTE) exists. This ratio is enhanced for most regions in NGC 3603; enhanced β/α ratios in other sources have been attributed to high optical depth or stimulated emission. We compare the morphology and kinematics of the ionized gas at 3.4 cm with the distribution of stars, $10\mu\text{m}$ emission and H₂O, OH, and CH₃OH maser emission. These comparisons suggest that both NGC 3576 and NGC 3603 have undergone sequential star formation.

Subject headings: HII regions: individual (NGC 3576, NGC 3603) – HII regions: kinematics and dynamics – ISM: abundances

1. Introduction

NGC 3576 ($D = 3.0 \pm 0.3 \text{ kpc}$) and NGC 3603 ($D = 6.1 \pm 0.6 \text{ kpc}$) are two of the highest luminosity optically visible HII regions in our Galaxy (Goss & Radhakrishnan, 1969). Indeed

NGC 3603 (with a bolometric luminosity of $10^7 L_{\odot}$) contains some $10^4 M_{\odot}$ of ionized gas, and aside from W49A, may be the most massive HII region in the Galaxy (Eisenhauer et al. 1998). Sources such as these are ideal laboratories to study the formation and evolution of massive stars and the initial mass function (IMF) of giant star forming regions. Similar regions that are visible with the Very Large Array (VLA) have been studied in detail. In particular, radio recombination line (RRL) studies of the Sgr B2 and W49A star forming regions with the VLA (Gaume et al. 1995; De Pree et al. 1995; Mehringer et al. 1995) have revealed small-scale outflows, very broad recombination lines ($\Delta V_{FWHM} \sim 80 \text{ km s}^{-1}$), expanding shells of ionized gas, and extremely compact structures. NGC 3576 and NGC 3603 appear to be similar regions in terms of luminosity, total mass and gas kinematics. Existing optical studies of NGC 3603, for example, have revealed broad emission lines and a stellar wind bubble, indicative of the early stages of a star forming region coming into equilibrium with its environment (Clayton 1986).

Though close in projection ($\sim 40'$), the two regions are separated by about 30 km s^{-1} in radial velocity as indicated by HI absorption (Goss & Radhakrishnan 1969) and hydrogen recombination line measurements (Wilson et al. 1970). Both regions are partially obscured by foreground dust, and studies of the kinematics of the associated ionized gas have been limited to optical wavelengths (Clayton 1986; Clayton 1990; Balick et al. 1980). Thus, existing velocity fields of the ionized gas are incomplete. The low resolution ($4'$) 6 cm image of Goss & Shaver (1970) shows that the radio emission associated with the two regions extends well beyond the detected $H\alpha$ emission. In both sources, however, it appears that the large scale ionized structures may be driven by energetic processes occurring near their cores. Evidence from a variety of wavelengths (including the radio line and continuum data from this study) indicate that both regions may have undergone sequential star formation, in NGC 3576 from east to west, and in NGC 3603 from north to south.

NGC 3603 and NGC 3576 have not been previously imaged with the Australia Telescope Compact Array (ATCA). Radio observations have been limited to low resolution line and continuum observations such as the $\sim 1'$ observations of Retallack & Goss (1980) at 21 cm. Thus, our radio frequency observations ($\theta_{FWHM} \sim 7''$) improve by factors of ~ 10 the best spatial resolution radio observations of the ionized gas associated with NGC 3576 and NGC 3603. For both NGC 3576 and NGC 3603, we present high resolution radio continuum images and radio recombination lines. New kinematic distances which are in reasonable agreement with the previously derived spectroscopic distances are derived from the observed line velocities and a revised Galactic radius of $R_0 = 8.5 \text{ kpc}$. We examine the ionized gas morphology and kinematics in the two sources, and possible star formation histories of these two regions. In addition, we derive electron temperatures, $H113\beta/H90\alpha$ ratios, helium abundances, and (where possible) carbon line strengths.

2. Observations and Data Reduction

Radio continuum and recombination line observations were made for NGC 3576 and NGC 3603 at the Australian Telescope Compact Array (ATCA) in the (750 m) B and (750 m) D configuration in two observing runs on the 13th and 19th of June in 1995 with a total of 24 hours of observing time. The observing frequency was 8.873 GHz, close to the frequency of the $H90\alpha$ recombination line. The wide bandwidth (16 MHz) allowed the simultaneous observation of three additional recombination lines: $He90\alpha$, $C90\alpha$, and $H113\beta$. Detailed observing parameters are presented in Table 1. The initial calibrations were made using the AIPS (Astronomical Image Processing System) software distributed by NRAO. Each data set was flagged and the flagged continuum data were then calibrated using the observed phase and flux density calibrators. These flags and calibrations were copied to the line data, where a bandpass calibration was applied and the continuum was subtracted from the u - v plane using the AIPS task UVLIN. Continuum images (with angular resolution of $\sim 7''$) were made from the line free channels for each source using the AIPS task IMAGR. This spatial resolution corresponds to a linear size of approximately 0.09 pc for NGC 3576 ($D = 3.0$ kpc) and approximately 0.21 pc for the more distant NGC 3603 ($D = 6.1$ kpc). The noise in the continuum images is dominated by the dynamic range limitations; thus the rms noise is significantly reduced in the continuum-subtracted line data.

After the initial data reduction in AIPS, the line and continuum data were analysed using GIPSY (the Groningen Image Processing System). Integrated line profiles were made for each pixel above a continuum cutoff level for both NGC 3576 ($5\sigma = 90$ mJy beam $^{-1}$) and NGC 3603 ($5\sigma = 55$ mJy beam $^{-1}$) using the GIPSY task PROFIL. Gaussian functions were then fitted to the spectral line data. The output from these fits are the line to continuum ratio (T_L/T_C), the integrated line strength, the central velocity (V_{LSR}) and the velocity full width at half maximum (ΔV_{FWHM}) of the line emission in each region. The HII region NGC 3576 consists of a single extended source (containing several local bright spots) with an angular size of $\sim 75''$ (~ 0.7 pc). In NGC 3603, thirteen individual sources were observed above a 5σ level in the continuum image.

3. Results

3.1. 3.4 cm Continuum Emission

Continuum images at 3.4 cm were made for both NGC 3576 and NGC 3603 using the AIPS task IMAGR. Figure 1 is the resulting 3.4 cm continuum image for NGC 3576. The crosses in the plot indicate the positions of the 10 μ m sources tabulated by Frogel & Persson (1974). Table 2 lists the continuum parameters for NGC 3576. These parameters include peak flux density (S_{peak}), integrated flux density (S_{tot}), angular diameter in arcseconds (θ), and linear radius in pc (r). Using the Parkes 64-m telescope, McGee & Newton (1981) report a 14.7 GHz flux density of 97.8 Jy in NGC 3576. We detect about 73% of the single dish flux density, or 71 ± 7 Jy. From the

13 sources in NGC 3603, we detect a total flux density of 25 Jy, or 38% of the reported single dish flux density of 65.9 Jy.

Figure 2 shows the 3.4 cm continuum image of NGC 3603. The crosses indicate the positions of the known 10 μm sources in NGC 3603 (Frogel et al. 1977). Each emission peak higher than the continuum 5σ cutoff (55 mJy beam^{-1}) has been designated as a separate region. The regions are labeled A to M, corresponding to increasing right ascension. The AIPS task JMFIT fitted two dimensional Gaussians to the individual continuum sources in NGC 3603. Table 2 presents continuum parameters for each region (A-M) in NGC 3603. These parameters include peak flux density (S_{peak}), integrated flux density (S_{tot}), deconvolved geometrical mean diameter in arcseconds (θ), and the linear radius in pc (r). Table 3 presents derived physical parameters for the regions in NGC 3603. These calculated parameters include electron density (n_e), emission measure (EM), mass of ionized gas (M_{HII}), excitation parameter (U), Lyman continuum photon flux (N_{LyC}) and continuum optical depth (τ_c). The continuum parameters were calculated assuming a uniform density, spherical, ionization bounded HII region, using the corrected formulae of Mezger & Henderson (1969). The calculated values are of course dependent on these assumptions, and in a complex region like NGC 3603, these assumptions may be overly simplistic. Figures 1 and 2 are plotted in B1950 coordinates. The last two figures in the paper (Figures 8 and 9) are plotted in J2000 coordinates to help in comparisons between the radio continuum emission and images at other frequencies, since images in the literature are shown in both epochs.

3.2. Radio Recombination Lines

3.2.1. NGC 3576

A continuum-weighted, integrated line profile was generated for NGC 3576 using the GIPSY task PROFIL. The integrated profile was generated for all pixels above a continuum 5σ cutoff level. In addition, integrated profiles were generated for each pixel above the continuum 5σ level in a $7'' \times 7''$ box centered on each of the positions of the Irs sources tabulated by Frogel & Persson (1974). The integrated source profile, and the four profiles line profiles were then fitted with Gaussian functions using the GIPSY task PROFIT. Figure 3a presents the integrated line profile and model fit for NGC 3576. Each of the four lines ($\text{H}90\alpha$, $\text{He}90\alpha$, $\text{C}90\alpha$ and $\text{H}113\beta$) is clearly detected, and labeled in Figure 3a. The integrated spectra for the Irs sources (Irs1/2, Irs3, Irs4 and Irs5; Frogel & Persson 1974) in NGC 3576 are shown in Figures 4a-d. The spectra in Fig. 3a and Figs. 4a-d show the data (points), Gaussian fits (solid line), and the residuals (dotted line) for each recombination line. The parameters of Gaussian fits to the lines (V_{LSR} , ΔV_{FWHM} , and T_l/T_c) are presented in Table 4.

From the integrated emission from NGC 3576, we determine a helium abundance of $Y^+ = 0.09 \pm 0.01$, and a $\text{H}113\beta/\text{H}90\alpha$ ratio close to the theoretically predicted value of 0.276 ($\beta/\alpha = 0.25 \pm 0.01$). The electron temperature in NGC 3576 (T_e^*) is calculated assuming local

thermodynamic equilibrium (LTE) from both the H90 α and H113 β recombination lines (using equation [22] from Roelfsema & Goss [1992]):

$$T_e^* = (6943\nu^{1.1} \frac{T_c}{T_l \Delta V_{FWHM}} \frac{1}{1 + Y^+})^{0.87} K, \quad (1)$$

with line width (ΔV_{FWHM}) given in km s⁻¹ and ν in GHz. These LTE temperatures are given in Table 4. An electron temperature (T_e) may also be calculated with corrections made for non-LTE effects (using Eq. [23] from Roelfsema & Goss [1992]):

$$T_e = T_e^* [b_n (1 - \frac{\beta_n T_c}{2})]^{0.87} K, \quad (2)$$

where b_n and β_n values are from the tables of Salem & Brocklehurst (1979).

For NGC 3576, a turbulent velocity (v_{turb}) and an electron temperature (T_e) are calculated by solving two equations for line width (Eq. 3) using the hydrogen and helium (H90 α and He90 α) recombination line widths. The line width can be expressed as

$$< v_{turb}^2 > = \frac{3\Delta V_H^2}{8\ln 2} - \frac{3kT_e}{M} \quad (3)$$

where the v_{turb} is the turbulent velocity in km s⁻¹, k is the Boltzmann constant, M is the mass of the atom, T_e is the electron temperature, and ΔV_H is the line FWHM in km s⁻¹. Using this method for NGC 3576, we derive a turbulent velocity $v_{turb}=17\pm1.0$ km s⁻¹ and an electron temperature of (T_e) of 6100 ± 200 K.

In order to examine the spatial variation in the H90 α line parameters, single Gaussian fits were made to each pixel above the continuum 5σ level in NGC 3576. The results of these point-to-point fits to the H90 α line are presented in Figures 5a and 5b. Figure 5a shows the spatial variation in the the central line velocity (V_{LSR}), with continuum contours overlaid on the greyscale representation of the ionized gas velocity. Figure 5b shows the spatial variation in the linewidth (ΔV_{FWHM}) of the H90 α line. For NGC 3576, a map of the LTE electron temperature was also generated, using Equation 1 (above) for each point above 10σ in the continuum image. The resulting LTE temperature image is shown in Figure 5c, indicating a clear E-W temperature gradient. This temperature gradient has also been plotted using the AIPS task IRING, which averages emission in concentric semicircles. The semicircles (with a thickness of $\sim 7''$), are centered on the continuum peak of NGC 3576 and extend to the east. A plot of temperature as a function of distance from the peak (in arcsec) is presented as Figure 6. The apparent temperature gradient is discussed in §4.2.1.

3.2.2. NGC 3603

An integrated line profile was generated for the NGC 3603 region above a 5σ cutoff in the continuum emission. Figure 3b presents the integrated line profile and model fit for NGC 3603.

Because of the large variation in line velocities across the source, the carbon line (apparent in some of the individual source profiles in NGC 3603) was fitted with the line width fixed at 10 km s^{-1} . Gaussian fits were made to the integrated profile, and the parameters of these fits are given in Table 5. Figures 7a-7m present the radio recombination line profiles and Gaussian fits for each of the thirteen subregions (A-M) in NGC 3603. For all 13 subregions, the signal-to-noise ratio is sufficiently high to fit the $\text{H}90\alpha$, $\text{He}90\alpha$, and $\text{H}113\beta$ lines. The $\text{C}90\alpha$ recombination line was successfully fit only in regions NGC 3603 B and H with a fixed carbon line width. Sources in which parameters were fixed and the fixed values are indicated parenthetically in Table 5. In a few cases, the fitted linewidth (ΔV_{FWHM}) of the $\text{He}90\alpha$ line may be broad due to blending with the $\text{C}90\alpha$ line in the sources where a $\text{C}90\alpha$ line was not separately fitted. The line parameters from the Gaussian fits to the integrated emission from NGC 3603 and its 13 subregions are presented in Table 5. As with NGC 3576, the table also includes the LTE electron temperature (T_e^*), the corrected electron temperature (T_e), the ionized helium to hydrogen ratio (Y^+), and the $\text{H}113\beta$ to $\text{H}90\alpha$ ratio (β/α), and the turbulent velocity (v_{turb}). For NGC 3603, the non-LTE electron temperature was calculated using Equation 2 (above). The turbulent velocity was calculated from the width of the $\text{H}90\alpha$ line and the non-LTE electron temperature using Eq. 3 (above). These temperatures and turbulent velocities are given in Table 5.

4. Discussion

4.1. Distances to NGC 3576 and NGC 3603

McGee & Newton (1981) detected recombination lines at 14.7 GHz from 25 high emission measure HII regions using the Parkes 64-m radio telescope. Both NGC 3576 (1109-610) and NGC 3603 (1112-610) were included in their survey. McGee & Newton detected the $\text{H}76\alpha$ line from NGC 3576 and NGC 3603 at $-24.1 \pm 0.1 \text{ km s}^{-1}$ and $7.7 \pm 0.1 \text{ km s}^{-1}$ respectively. In addition to $\text{H}76\alpha$, McGee & Newton detected the $\text{He}76\alpha$ line toward both sources, and the carbon line toward NGC 3576. For NGC 3576, the detected line width in the $\text{H}90\alpha$ line ($\Delta V_{FWHM} = 28.7 \pm 0.1$) is in good agreement with the $\text{H}76\alpha$ line width published by McGee & Newton ($\Delta V_{FWHM} = 29.1 \pm 0.1$). The integrated $\text{H}90\alpha$ line width for NGC 3603 ($\Delta V_{FWHM} = 36.9 \pm 0.25$) is in reasonable agreement with the value for the $\text{H}76\alpha$ line given by McGee & Newton ($\Delta V_{FWHM} = 41.9 \pm 0.2$).

The distance to NGC 3576 has been discussed at length in two papers, Goss & Radhakrishnan (1969), and Persi et al (1994). Goss & Radhakrishnan found a kinematic distance of 3.6 kpc, just beyond the tangential point of the Galaxy, using the Schmidt galactic model and a value of $R_0 = 10 \text{ kpc}$. This distance is derived from a line center velocity of -23 km s^{-1} , and circular motion within the galaxy. This kinematic distance agrees well with the findings from Caswell & Haynes (1987). Our new observations, using a revised $R_0 = 8.5 \text{ kpc}$, and a line velocity of $-20.0 \pm 0.1 \text{ km s}^{-1}$ suggest a distance of $3.0 \pm 0.3 \text{ kpc}$ for NGC 3576. Because no ionizing cluster is optically visible, no reliable spectroscopic distance has been determined. Humphreys (1972), in a study of

the effect of streaming in the Carina arm, determined a distance of 3.2 kpc for NGC 3576, and detected both velocity variations ($\sim 4 \text{ km s}^{-1}$) and a large spread in the individual velocities of early type stars in this direction. Persi et al. (1994) derive a distance based on the star HD 97499 (position plotted in Figure 8). However, the high resolution radio continuum image of the source (Fig. 8) shows that there may not be a direct association between HD 97499 and NGC 3576. HD 97499 (RA = 11 12 10.1, Dec = -61 18 45.1 J2000) is located $\sim 2'$ east of the peak in the 3.4 cm continuum emission. As a result, it is unlikely that HD 97499 contributes greatly to the ionization seen in NGC 3576. Finally, the stellar velocity of HD 97499 ($-32 \text{ km s}^{-1} \pm 11 \text{ km s}^{-1}$; Crampton, 1972) does not agree well with the observed velocities in the ionized gas (-23 km s^{-1} [McGee & Gardner, 1968] and -25 km s^{-1} [Caswell & Haynes, 1987]). Thus, distances associated with HD 97499 should not simply be applied to NGC 3576.

The determination of the distance to NGC 3603 is aided by the presence of an optically visible ionizing cluster. Distance estimates made from stellar observations can be compared to kinematic distances derived from the ionized gas. The earliest spectroscopic distance estimate was made by Sher (1965) at 3.5 kpc. Without a large number of faint stars visible, the spectroscopic distance estimate was admittedly crude. Shortly thereafter, Goss & Radhakrishnan (1969) found a distance of 8.4 kpc based on the observed line velocity of 10 km s^{-1} , using $R_0 = 10 \text{ kpc}$. Caswell & Haynes (1987) also found a kinematic distance using $R_0 = 10.5 \text{ kpc}$ and $V = +11 \text{ km s}^{-1}$ of 8.6 kpc. McGee & Newton (1982) report an H76 α line velocity of $7.7 \pm 0.1 \text{ km s}^{-1}$. Our velocity for the integrated line in NGC 3603 is $V_{LSR} = 9.1 \pm 0.1$, consistent with previously reported values. The H90 α velocities found for the thirteen individual regions range between -2 km s^{-1} and 19 km s^{-1} . Using the velocity from the integrated line ($V_{LSR} = 9.1 \text{ km s}^{-1}$) and $R_0 = 8.5 \text{ kpc}$, we determine a kinematic distance of $6.1 \pm 0.6 \text{ kpc}$.

UBV stellar photometry of NGC 3603 has resulted in distance estimates as low as 3.5 kpc (Sher, 1965) and as high as 8.1 kpc (Moffat, 1974). Melnick & Grosbøl (1982) derive a distance of 5.3 kpc. Moffat (1974), using the stars observed by Sher (1965) and a different reddening scale derive a distance of $8.1 \pm 0.8 \text{ kpc}$. Finally, Melnick, Tapia & Terlevich (1989) use *UBV* photometry of the ionizing cluster of NGC 3603 to find a distance modulus of $(m - M)_0 = 14.3$, which corresponds to a distance of $\sim 7.2 \text{ kpc}$. Their value is in good agreement with both Moffat (1974), and the kinematic distances of Goss & Radhakrishnan (1969), and Caswell & Haynes (1987), and in reasonable agreement with the distance derived from our integrated radio recombination line velocity. The smaller distances inferred from the measurements of Sher (1965) and Melnick & Grosbøl (1982) are likely the result of errors in photometry due to nebular contamination (Melnick, Tapia & Terlevich 1989).

4.2. NGC 3576

NGC 3576 has been extensively observed in the radio continuum (Goss & Shaver 1970) and in radio recombination lines (McGee & Gardner 1968, McGee & Newton 1981, Wilson et al. 1970).

Recombination line studies of the region show the line velocities to be centered at approximately -23 km s^{-1} . The core (diameter $\sim 1.5'$) of NGC 3576 is the location of five identified infrared ($10\mu\text{m}$) sources (Frogel & Persson 1974), as well as CH_3OH (Caswell et al. 1995) and H_2O (Caswell et al. 1989) masers. The presence of bright $10\mu\text{m}$ emission, water masers and compact thermal radio emission are typical indications of the early stages of star formation and dense circumstellar environments.

Persi et al. (1994) have identified 135 individual stars with IR excess (to distinguish cluster stars from red field stars) in an extremely young galactic cluster associated with the HII region NGC 3576, using *JHK* images and photometry. Approximately 40 of these IR sources are members of the cluster. In their *JHK* images, Persi et al. (1994) detect a color gradient in the near-IR, indicating increasing extinction toward the western edge of NGC 3576. This gradient suggests that star formation may have progressed in NGC 3576 from east to west, with the youngest stars currently located at the western edge of the source.

In addition, Persi et al. (1994) provide a summary of existing studies of the region, and an overlay of the 21 cm continuum (Retallack & Goss 1980), the $10 \mu\text{m}$ continuum sources (Frogel & Persson 1974) and their K band mosaic of the region. The overlay (their Fig. 7) clearly shows that the radio continuum, near-IR continuum and $10\mu\text{m}$ continuum are all strongly peaked toward the western edge of the source. Our 3.4 cm radio continuum image (Fig. 1) improves significantly on the resolution of Retallack & Goss (1980), and shows that there is a coincidence between the location of the infrared sources (Irs1-Irs5) tabulated by Frogel & Persson (1974) and the extended ionized radio continuum emission. Figure 1 shows the 3.4 cm radio continuum with crosses and labels indicating the positions of the $10\mu\text{m}$ peaks (Frogel et al. 1977). All five sources are coincident with the ionized gas emission and are near local peaks in the radio continuum emission.

4.2.1. Characteristics of the Ionized Gas

Our radio recombination line observations have allowed us to examine in detail the variation in electron temperature and helium abundance in NGC 3576. Our results in both cases seem to strengthen the case for sequential star formation in that source. The LTE electron temperatures show a clear gradient from east to west, with temperature increasing with distance (east) from the continuum peak. The gradient is apparent in Figures 5c and 6. Figure 5c shows the spatial variation in electron temperature across the source. Figure 6 is a plot of temperature as a function of distance from the 3.4 cm continuum peak. The LTE electron temperature increases with distance from 6000 K to a maximum of $\sim 8,000$ K, and then falls to lower levels in the more diffuse gas to the east. If the ionizing stars are those IR sources coincident with the radio continuum, this rising temperature signature is expected (Hjellming 1966). Lower energy photons on average would be absorbed first, with the higher energy photons penetrating deeper into the diffuse gas. Thus, the temperature gradient confirms a scenario in which the youngest, hottest stars in the HII region are located near the sharp western edge of the radio continuum.

Under LTE conditions, the theoretical value of the H113 β to H90 α hydrogen ratio is 0.276 (Shaver & Wilson, 1979). Deviations from this theoretical ratio indicate local deviations from LTE, and the ratio can be used to correct the derived electron temperature. The H113 β to H90 α ratio in NGC 3576 varies from $\sim 0.25 \pm 0.05$ to $\sim 0.35 \pm 0.05$, with the lowest values associated with the bright peak in the radio continuum along the western edge of the HII region. Pressure broadening can lower the β/α ratio from the theoretical value. When the β/α ratio is lower than the theoretical value, electron temperatures derived from the β lines will of course be higher than those derived from α lines. In fact, for NGC 3576, the electron temperatures derived from the H113 β line are ~ 11 % higher than those derived from the H90 α recombination line.

A small number of Galactic objects are known to have helium abundances in excess of 20%. Such elevated helium abundances have been reported in the W3 core (Adler, Wood & Goss 1996, Roelfsema, Goss & Mallik 1992), and are thought to be associated with the environments of evolved stellar objects that are rapidly losing mass. A small but significant positive gradient in helium abundance (Y^+) is detected from the western peak of the source ($Y^+ = 0.08 \pm 0.01$) to the more diffuse ionized gas to the east ($Y^+ = 0.14 \pm 0.03$). This range is consistent with the Y^+ value reported by McGee & Newton (1981) of $N(\text{He}^+)/N(\text{H}^+) = 0.091$. The abundance gradient is consistent with the suggestion of Persi et al. (1994) who detect a color gradient in the near-IR indicating increasing extinction to the west of the source. Persi et al. (1994) propose that the youngest stars are located at the western edge of the ionized gas. The higher helium abundances found to the east of NGC 3576 may indicate the presence of a population of older stars and their associated mass loss.

4.2.2. Gas Kinematics in NGC 3576

Figures 5a and 5b show that both broad lines and velocity gradients are present in the ionized gas associated with NGC 3576. Double-peaked emission lines are located over a $\sim 15''$ region at the northern edge of the source. In Fig. 5a, the double-peaked lines have been fitted with a single Gaussian with a $\Delta V_{FWHM} = 50 \text{ km s}^{-1}$. Closer inspection (see Figure 1 inset) shows that locally the lines are possibly double-peaked. The double-peaked recombination lines in this region have a velocity separation of $\sim 25 \text{ km s}^{-1}$, and may be due to either the expansion of an ionized shell of gas, or the outflow of ionized gas from a local source. The broad line region is located to the east of the Irs1/Irs2 peak (Frogel & Persson 1974). The region of the broadest radio recombination lines corresponds spatially to a distinct gap in the $10 \mu\text{m}$ emission between Irs1/Irs2 and Irs5.

If the split lines are due to expansion, then the rate of expansion is $\sim 13 \text{ km s}^{-1}$. Double-peaked lines with similar velocity separations have been seen in some of the compact HII regions in Sgr B2 North (De Pree et al. 1996). The spatial resolution of the current data is insufficient to determine whether an expanding shell of ionized gas is located at this position. Alternatively, the double-peaked lines may arise near the base of a large scale ionized outflow.

A large, approximately north-south velocity gradient of $30 \text{ km s}^{-1} \text{ pc}^{-1}$ is observed in the ionized gas, and is shown in Figure 5a. The velocity across the source varies from $\sim 0 \text{ km s}^{-1}$ along the southern edge to $\sim 35 \text{ km s}^{-1}$ in the north. The gradient is not likely due to rotation in the ionized gas, since these rotational velocities would imply an approximate enclosed mass of $M = 2 \times 10^5 M_{\odot}$. The gradient is more likely due to an ionized outflow, and may in fact be the base of a much larger scale outflow. A wide field of view optical image of the region (Fig. 8; Persi et al. 1994) shows several large loops and filaments of ionized gas extending several minutes of arc to the north of NGC 3576.

4.2.3. *Sequential Star Formation in NGC 3576*

Figure 8 shows the positions of cluster members with infrared excess (crosses; Persi et al. 1994) overlaid on the 3.4 cm radio continuum. Approximately 40 infrared excess sources have been identified. Eight of the stellar positions are coincident with the radio continuum emission, with the majority of the sources located to the north, east and south of the radio continuum. Only a single infrared excess source is located to the west of the radio continuum peak. This spatial distribution would be expected if star formation were proceeding from east to west into the molecular cloud. Water (H_2O) and methanol (CH_3OH) maser emission (plotted) are located coincident with the peak in the radio continuum emission.

The location of molecular gas in NGC 3576 is also consistent with a triggered star formation scenario. The peak in the 1.0 mm dust continuum emission (Cheung et al. 1980) is located to the west of the bright radio continuum peak. The 1.0 mm emission traces the core of the molecular cloud located to the west of NGC 3576. Similar distributions of ionized and molecular gas are found near the Sgr B2 F HII regions (De Pree et al. 1996), G34.3 (Gaume et al. 1994), and others. In fact, the morphology of NGC 3576 is “cometary”, and closely resembles a mirror image of the cometary source G34.3 (Gaume et al. 1994) in which a dense molecular cloud core is located at the eastern periphery of the ionized gas.

Taken as a whole, the radio, infrared, optical and molecular imaging of NGC 3576 suggest that the oldest ionizing sources are located to the east of the region, and the youngest ionizing sources to the west. The distribution of stars, ionized gas, molecular gas, and water masers in NGC 3576 is reminiscent of simple scenarios of sequential star formation. In this picture, the star formation process slowly “eats away” at a dense cloud, so that the most evolved stars are located far from the cloud core. The ionized gas is found at the periphery of the cloud, and the youngest sources (protostars) are still embedded in the nascent molecular material, but reveal their presence with water maser emission. In many such regions in the Galaxy, the stellar population is not observable. The ability to observe stars, ionized gas and molecular gas in this distant source make it unique in the Galaxy.

4.3. NGC 3603

NGC 3603 fills a gap in luminosity between a region like Orion, with ~ 1 O-type star, and R136 in 30 Doradus, with ~ 100 O-type stars (Eisenhauer et al. 1998). The ionizing core of NGC 3603 (HD 97950) consists of approximately 20 O-type and Wolf-Rayet stars (Moffat, 1983). From photometric measurements, Melnick, Tapia & Terlevich (1989) have identified 181 sources (50 with U , B , and V photometry) associated with the stellar cluster within NGC 3603. The authors found an average age of 2.5 ± 2 Myr and evidence that star formation propagated from north to south, ending at the bordering molecular cloud (Cheung 1980). Brandner et al. (1997) have identified a ring-shaped nebula and bipolar outflows from Sher 25, a cluster member of HD 97950. Sher 25 (a B1.5Iab supergiant) is located approximately $\sim 20''$ north of the core of the cluster. Brandner et al. present evidence for association of this star with HD 97950 and compare the star to the progenitor of SN 1987A.

The initial mass function (IMF) of HD 97950, the stellar cluster associated with NGC 3603, has been discussed in recent articles by Moffat et al. (1994), Hofmann et al. (1995), Zinnecker (1995) and Eisenhauer et al. (1998). Because of the proximity of NGC 3603, high resolution, deep studies allow detailed counts of stellar mass distribution to be made. Moffat et al. (1994) observed the central cluster with the HST/PC, studying its dense stellar core and the population of Wolf-Rayet stars. Moffat et al. (1994) conclude that in terms of density and stellar population, NGC 3603 is a ‘Galactic clone’ of R136 in 30 Doradus. Hofmann et al. derive an IMF power law of $\Gamma = -1.4 \pm 0.6$ ($\xi \propto M^\Gamma$) for 30-60 $M_\odot$ which is similar to that of 30 Doradus. Using speckle masking techniques with the HST, Hofmann et al. (1995) found a similar slope for stellar masses of 15 to 50 $M_\odot$ of $\Gamma = -1.6 \pm 0.2$. Both Eisenhauer et al. and Zinnecker use adaptive optics to extend the IMF to stellar masses less than 1 $M_\odot$. Some theoretical and observational studies have claimed that starburst systems are deficient in low mass stars, but Eisenhauer et al. (1998) find that the IMF follows a Salpeter power law with an index of ≤ -0.73 , and that there does not appear to be a low mass truncation to the IMF down to the observational limit of 0.2 $M_\odot$. From such properties of NGC 3603, it may be possible to extrapolate properties of larger extragalactic starbursts.

Sequential star formation has also been suggested in NGC 3603, proceeding from north to south (Melnick, Tapia & Terlevich 1989). The presence of evolved objects at the north end of the source and embedded IR sources and a molecular cloud to the south qualitatively confirms a sequential star formation scenario (Hofmann et al. 1995). Investigations of NGC 3603 have proceeded on two fronts: radio and optical observations of the ionized gas, and optical and infrared observations of the exciting stars. While the stars and gas in the giant HII region are visually observable, these high resolution radio observations present the first unobscured view of the ionized gas on small scales. Melnick, Tapia & Terlevich (1989) have tabulated the positions of 181 stars in NGC 3603 from their CCD UBV photometry. The positions of the 50 cluster member stars for which U , B , and V magnitudes are tabulated are shown in Figure 9. The figure shows the positions of these stars (crosses) plotted with the 3.4 cm radio continuum. Coordinates for

the stellar positions were derived from pixel coordinates using the known position of the B1.5Iab supergiant (Sher 25) as given by Brandner et al. (1997).

Optical studies indicate both the presence of high velocity gas and of highly evolved stars. Clayton (1986) first investigated the velocity structure in the ionized gas located at the core of NGC 3603, and found both large scale motions (perhaps kinematic evidence of earlier supernovae) and two smaller shells, possibly wind-driven bubbles. The $H\alpha$ and [NII] line profiles were made along two lines across the core of NGC 3603. One possible explanation of the wind-driven bubbles was suggested by Moffat et al. (1994) who detected three Wolf-Rayet (WR) stars in their HST/PC observations of the cluster core. Hofmann et al. (1995) confirm the presence of three, perhaps four WR stars in the cluster core. Brandner et al. (1997) provide fascinating evidence that one of the sources believed to be an older member of the cluster (Sher 25, a B1.5Ia supergiant) is at the center of a clumpy ring of optical emission and a possible bipolar outflow. They compare the source to the progenitor of SN1987A.

Caswell & Haynes (1987) detected strong hydrogen radio recombination line emission from NGC 3603. The source was also included in the recombination line survey of McGee & Newton (1981), and both hydrogen and helium (at 14.7 GHz) were detected. Our integrated $H90\alpha$ line width ($\Delta V_{FWHM}=36.9\pm0.25$) is in reasonable agreement with the value for the $H76\alpha$ line given by McGee & Newton ($\Delta V_{FWHM}=41.9\pm0.2$).

The present study of the ionized gas in NGC 3603 addresses a number of the outstanding issues in this source:

1. Are there any regions of enhanced helium abundance, as might be expected in the presence of 3-4 WR stars?
2. Is there any evidence in the ionized gas emission of the bipolar outflow associated with the B1.5 supergiant Sher 25 (Brandner et al. 1997)?
3. How does the location of ionized gas compare to the known positions of infrared sources (Frogel et al. 1977), WR stars (Moffat et al. 1994 and references therein) and maser emission (Caswell et al. 1989, Caswell et al. 1998)?

4.3.1. Helium Abundance and β/α Ratios

McGee & Newton (1981) derived a helium abundance in NGC 3603 of $Y^+ = 0.069$ (no uncertainty given). Our calculated helium abundance from the integrated line profile is significantly higher ($Y^+ = 0.13\pm0.01$), but in their notes on individual sources, McGee & Newton (1981) indicate that their $He76\alpha$ line profile was rather noisy.

Moffat et al. (1994) detect three prominent Wolf-Rayet stars in NGC 3603 in their narrowband HeII (4686 Å) images. WR stars have been cited as the possible explanation of enhanced helium

abundances (Adler, Wood & Goss 1996). It is certainly reasonable to look for evidence of such enhanced abundances in the vicinity of known WR stars. In fact, the highest helium abundances in NGC 3603 are detected in source C, which has a value of $Y^+ = 0.26 \pm 0.10$. Interpretation of this high value is complicated by the large uncertainty. The ionized gas in NGC 3603 is located on the periphery of the WR wind-blown bubble, so it is possible that ionized helium (Y^+) abundances are enhanced by the presence of an evolved star.

The observed $H113\beta/H90\alpha$ ratios for the 13 sources in NGC 3603 range from 0.29 ± 0.04 to 0.53 ± 0.12 , all above the theoretical value ($\beta/\alpha=0.276$). Most detections are within 2σ of this theoretical value, but a few sources (NGC 3603 E, H, I, and K) exceed the theoretical value by more than 3σ . Anomalously high β/α ratios were found from $H92\alpha$ and $H115\beta$ recombination line observations in two Galactic center regions (G0.18-0.04; the Sickle and the Pistol; Lang, et al., 1997). Thum et al. (1995) propose a model that can explain enhanced β/α ratios as the result of high continuum optical depth. However, NGC 3603 A-M appear to have high β/α ratios, and yet be low density ($n_e \leq 10^4 \text{ cm}^{-3}$) regions with $\tau_c \leq 1$. Cersosimo & Magnani (1990) suggest that under such conditions, high β/α ratios may result from stimulated emission caused by a background source.

4.3.2. Ionized Gas Kinematics

No ionized gas is detected above a 5σ continuum level at the location of Sher 25. The optical study of Clayton (1986) indicates clearly that there are both large scale and small scale motions in the ionized gas. Plate 1 of Clayton shows an $H\alpha$ image of the central $6'$ of NGC 3603, and the positions of the two slit positions through the core. The bright ionized gas observed in our high resolution image (see Fig. 1b and Fig. 9) lies to the south of the cluster core. On scales probed by our high resolution observations, only one source (NGC 3603 F) has clearly double-peaked recombination line emission. Two Gaussians with velocities separated by $\sim 29 \text{ km s}^{-1}$ ($V_{LSR} = 27.5 \text{ km s}^{-1}$ and -1.1 km s^{-1}) have been fitted to the $H90\alpha$ line in NGC 3603 F. The velocity separation is similar to that seen in other Galactic wind-driven bubbles and also in NGC 3576. In the radio continuum, source F is a relatively faint, marginally resolved source located near the western edge of the southern bright region. NGC 3603 F has a deconvolved diameter of $\theta_{FWHM} \sim 25''$, giving the source a linear diameter of 0.5 pc ($D = 6.1 \text{ kpc}$). Thus, source F may be an expanding shell of ionized gas. A small optically observable wind-driven "stellar bubble" with $r \sim 0.6 \text{ pc}$ has been observed centered on the HD97950 star cluster in NGC 3603 (Balick, Boeshaar & Gull, 1980).

In NGC 3603, the individual source velocities (as derived from the $H90\alpha$ line) range from $-3.7 \pm 0.3 \text{ km s}^{-1}$ in source A to $15.9 \pm 0.2 \text{ km s}^{-1}$ in source I. The broadest lines are detected in NGC 3603 F ($\Delta V_{FWHM} = 52.7 \pm 2.2 \text{ km s}^{-1}$). As indicated above, the broad lines in this source likely result from the double-peaked lines apparent in Fig. 7f. Broad lines can result from a variety of local conditions including pressure broadening or spatially unresolved velocity gradients. Given

the relatively high frequency of the observations and other evidence for velocity gradients in the source, it is likely that the broad lines in source M (located to the north of the cluster core) are related to gas kinematics. NGC 3603 M has $v_{turb} \simeq 26 \pm 10 \text{ km s}^{-1}$. Other localized regions in the source have very broad recombination line emission. Lines as broad as $\Delta V \sim 80 \text{ km s}^{-1}$ are detected in the low surface brightness gas between sources NGC 3603 D and F. And in the bright core of NGC 3603 D ($\sim 0.07 \text{ pc}$ in diameter) line widths approach $\Delta V \sim 75 \text{ km s}^{-1}$. In a sequential star formation scenario, sources D and F are located at the boundary of the molecular cloud and the stellar cluster, and would be in the early stages of their evolution, perhaps still coming into equilibrium with their environment or photoevaporating material from the molecular cloud interface.

4.3.3. Evidence for sequential star formation in NGC 3603

In Figure 2, the positions of ten $10\mu\text{m}$ sources (Frogel et al. 1977) are plotted over the 3.4 cm continuum. Figure 9 shows the positions of cluster stars and known maser emission. Error bars for the absolute positions of the IR sources are $\pm 10''$, and the size of the crosses in Fig. 2 indicates the absolute error in the positions of the infrared sources. In several cases (e.g. Irs 1, 2, 9, 14), the infrared sources are coincident with radio continuum sources.

Perhaps the clearest evidence for sequential star formation in NGC 3603 is the relative location of the radio emission from the ionized gas, the stellar positions, and the position of embedded infrared sources and maser emission. The distribution is very much like that observed in NGC 3576. The most evolved stars are located to the north, centered in what appears at many frequencies to be the wind-blown bubble resulting from the WR stars located there. The molecular cloud core is located farthest to the south, and the ionized gas emission (apparent at optical and radio frequencies) is located at the periphery between the stars and the cold gas.

5. Conclusions

Both NGC 3576 and NGC 3603 provide rare opportunities to study massive star formation in a variety of spectral regions. Massive star forming regions like W49A and Sgr B2 are impressive at radio frequencies, but the stellar populations are inaccessible in the infrared. In these two sources, high resolution stellar photometry and radio continuum and recombination line observations provide a more complete picture of the physical conditions and complex gas kinematics.

The following is a summary of our major conclusions:

- (1) In these high resolution radio continuum and recombination line observations, we have identified a number of previously unresolved radio continuum sources, particularly in NGC 3603.
- (2) The observed temperature gradient in NGC 3576 is in agreement with the known color

gradient in the source and supports a sequential star formation scenario that has been proposed for the region.

(3) We have detected several compact sources in NGC 3603 with broad recombination lines. One source (NGC 3603 F) has double peaked profiles, with the peaks separated by $\sim 30 \text{ km s}^{-1}$.

(4) The brightest radio continuum emission in NGC 3603 is located to the south of the core of the stellar cluster HD 97950. There is no radio continuum or line emission apparent at the position of Sher 25.

(5) Derived helium abundances and electron temperatures in NGC 3576 are in good agreement with values given in McGee & Newton (1981) while those in NGC 3603 are significantly higher.

(6) Enhanced helium abundances ranging in value from $Y^+ = 0.16$ to 0.26 are detected in NGC 3603 A-D and J. These regions may have had their helium abundance enriched by the 3 or 4 known WR sources located at the cluster core.

(7) In both sources, the relative positions of stars, ionized gas, molecular gas and masers is suggestive of a sequential star formation scenario. In NGC 3576 the star formation appears to have proceeded from east to west, in NGC 3603 from north to south.

C. G. De Pree gratefully acknowledges the support of an American Astronomical Society Small Research Grant. W. M. Goss would like to thank R. D. Ekers, J.B. Whiteoak, & D. McConnell for support during a sabbatical visit to the Australia Telescope Compact Array in 1996. The Australia Telescope is funded by the Commonwealth of Australia for operation as a National Facility operated by CSIRO. The National Radio Astronomy Observatory is a facility of the National Science Foundation, operated under cooperative agreement by Associated Universities, Inc. The authors also thank H. Zinnecker, J. Caswell, W. Brandner, P. Palmer, D. Mehringer, L. Rodriguez, R. Gaume, M. Claussen and an anonymous referee for helpful discussions and comments.

REFERENCES

- Adler, D.S., Wood, D.O.S., Goss, W.M. 1996, ApJ, **471**, 871
- Balick, B., Boeshaar, G.O., Gull, T.R. 1980, ApJ, **242**, 584
- Brandner, W., Grebel, E.K., Chu Y., Weis, K. 1997, ApJ, **475**, L45
- Caswell, J.L., 1998, MNRAS, 297, 215
- Caswell, J.L., Vaile, R.A., Ellingsen, S.P., Whiteoak, J.B., Norris, R.P. 1995, MNRAS, 272, 96
- Caswell, J.L., Haynes, R.F. 1987, A&A, **171**, 261
- Caswell, J.L., Batchelor, R.A., Forster, J.R., Wellington, K.J. 1989, *Austral. J. Phys.*, **42**, 331
- Caswell, J.L., Vaile, R.A., Ellingsen, S.P., Whiteoak, J.B., Norris, R.P., 1977, MNRAS, **272**, 96
- Cersosimo, J.C., Magnani, L. 1990, A&A, **239**, 287
- Cheung, L. H., Frogel, J. A., Hauser, M. G. & Gezari, D. Y. 1980, ApJ, 240, 74
- Clayton, C.A. 1986, MNRAS, **219**, 895
- Clayton, C.A. 1990, MNRAS, **246**, 712
- Crampton, D. 1972, MNRAS, **158**, 85
- De Pree, C.G., Mehringer, D. M. & Goss, W. M. 1997, ApJ, 482, 307
- De Pree, C.G., Gaume, R. A., Goss, W. M. & Claussen, M.J., 1996, ApJ, 464, 788
- De Pree, C.G., Gaume, R.A., Goss, W.M. & Claussen, M.J., 1995, ApJ, 451, 284
- Eisenhauer, F., Quirrenbach, A., Zinnecker, H., and Genzel, R. 1998, ApJ, 498, 278
- Frogel, J.A., Persson, S.E., Aaronson, M. 1977, ApJ, 213, 723
- Frogel, J.A., Persson, S.E. 1974, ApJ, **192**, 351
- Gaume, R.A., Fey, A.L. & Claussen, M.J. 1994, ApJ, 432, 648
- Gaume, R.A., Claussen, M.J., De Pree, C.G., Goss, W.M. & Mehringer, D.M. 1995, ApJ, 449, 663
- Goss, W.M., Radhakrishnan, V. 1969, Astrophys Letters, **4**, 199
- Goss & Shaver 1970, *Austral. J. Phys. Astroph. Suppl.*, 14, 133
- Hjellming, R. 1966, ApJ, 143, 420
- Hofmann, K.H., Seggewiss, W., Weigelt, G. 1995, A&A, **300**, 403
- Humphreys, R.A. 1972, A&A, **20**, 29
- Lang, C., Goss, W.M., Wood, D.O.S. 1997, ApJ, **474**, 275
- Mehringer, D.M., De Pree, C.G., Gaume, R.A., Goss, W.M. & Claussen, M.J. 1995, ApJ, 442, L29
- Mezger, P.G. & Henderson, A.P., ApJ, 147, 471
- McGee, R.X., Gardner, F.F. 1968, *Austral. J. Phys.*, **21**, 149

- McGee, R.X., Newton, L.M. 1981, MNRAS, **196**, 889
- Melnick, J., Grosbøl, P. 1982, A&A. **107**, 23
- Melnick, J., Tapia, M., Terlevich, R. 1989, A&A, **213**, 89
- Moffat, A.F.J., 1974, A&A, **35**, 315
- Moffat, A.F.J., 1983, A&A, **124**, 273
- Moffat, A.F.J., Drissen, L., Shara, M.M. 1994, ApJ, **436**, 183
- Norris, R. P., Caswell, J. L., Gardner, F. F., & Wellington, K. J. 1987, ApJ, 321, 159
- Persi, P., Roth, M., Tapia, M., Ferrari-Toniolo, M., Marenzi, A.R. 1994, A&A, **282**, 474
- Retallack, D.S., Goss, W.M., 1980, MNRAS, **193**, 261
- Roelfsema, P.R., Goss, W.M., 1992, A&A Review, **4**, 161
- Roelfsema, P.R., Goss, W.M., & Mallick, 1992
- Salem, M., & Brocklehurst, M. 1979, ApJS, 39, 633
- Shaver, P.A., Wilson, T.L., 1979, A&A **79**, 312
- Sher, D. 1965, MNRAS, **129**, 17
- Thum, C., Strelinski, V.S., Martin-Pintado, J., Matthews, H.E., Smith, H.A. 1995, A&A, **300**, 643
- Wilson, T. L., Mezger, P. G., Gardner, F. F., Milne, D. K. 1970, A&A, 6, 364
- Zinnecker, H., 1995, 11th IAP Conference (Paris)

Table 1: Observational Parameters of NGC 3576 and NGC 3603

Parameter	NGC 3576	NGC 3603
Date (configuration).....	13 June 1995 (750m D) 19 June 1995 (750m B)	13 June 1995 (750m D) 19 June 1995 (750m B)
Total observing time (hr).....	12	12
R.A. of field center (J2000).....	$11^h 11^m 51.88^s$	$11^h 15^m 02.6^s$
Decl. of field center (J2000).....	$-61^\circ 18' 31''$	$-61^\circ 15' 46''$
Major axis ($''$).....	$7.0''$	$7.0''$
Minor axis ($''$).....	$6.8''$	$6.9''$
Position angle ($^\circ$).....	-9°	-13°
LSR central velocity (km s^{-1}).....	-23	10
Total bandwidth (MHz).....	16	16
Number of channels.....	256	256
Channel Separation (kHz, km s^{-1})...	62.5 (2.1)	62.5 (2.1)
Spectral resolution (kHz, km s^{-1})...	75.6 (2.6)	75.6 (2.6)
Continuum noise (mJy beam^{-1}).....	18	11
Line noise (mJy beam^{-1}).....	2.1	2.2

Table 2: Continuum Parameters for NGC 3576 and NGC 3603

Source	RA (J2000)			Dec (J2000)			S_{Peak} (Jy beam ⁻¹)	$S_{Tot.}$ (Jy)	θ (")	r (pc)
NGC 3576	11 ^h	11 ^m	51.32 ^s	-61 ^o	18'	43''5	4.1±0.4	71±7	~90	1.4
NGC 3603 A	11 ^h	14 ^m	56.02 ^s	-61 ^o	13'	55''8	0.10±0.01	1.3±0.3	24	0.52
B		14 ^m	56.51 ^s		13'	17''0	0.08±0.01	2.5±0.4	38	0.80
C		15 ^m	00.90 ^s		13'	32''7	0.06±0.01	0.85±0.3	24	0.52
D		15 ^m	01.99 ^s		16'	33''0	0.11±0.01	1.2±0.3	22	0.46
E		15 ^m	02.62 ^s		15'	53''8	0.40±0.02	1.9±0.2	13	0.28
F		15 ^m	06.13 ^s		16'	40''5	0.12±0.01	1.7±0.3	25	0.52
G		15 ^m	08.76 ^s		16'	55''7	0.30±0.01	4.4±0.3	26	0.54
H		15 ^m	09.45 ^s		16'	41''4	0.23±0.01	4.6±0.3	30	0.64
I		15 ^m	10.36 ^s		16'	17''4	0.25±0.02	1.3±0.2	14	0.30
J		15 ^m	14.24 ^s		17'	34''0	0.11±0.01	2.2±0.3	30	0.65
K		15 ^m	18.75 ^s		16'	58''1	0.08±0.01	1.2±0.3	25	0.53
L		15 ^m	24.20 ^s		12'	54''0	0.06±0.01	0.94±0.3	29	0.61
M		15 ^m	31.59 ^s		13'	16''7	0.06±0.01	1.1±0.3	26	0.55

Table 3: Derived Continuum Parameters for NGC 3603

Source	n_e (10^3 cm^{-3})	EM (10^6 pc cm^{-6})	M_{HII} ($M_\odot$)	U (pc cm^{-2})	N_{LyC} (10^{48} s^{-1})	τ_c
A	0.92	0.9	13	50	5.6	0.005
B	0.67	0.7	35	62	11	0.004
C	0.75	0.6	10	44	3.7	0.005
D	1.0	1.0	10	49	5.1	0.006
E	2.8	4.6	6	57	8.2	0.027
F	1.0	1.1	15	54	7.1	0.007
G	1.6	2.8	25	76	19	0.016
H	1.3	2.1	32	76	20	0.012
I	2.1	2.7	5	50	5.4	0.016
J	0.87	1.0	23	60	9.5	0.006
K	0.85	0.8	12	48	5.0	0.005
L	0.66	0.5	15	47	4.8	0.003
M	0.72	0.6	12	45	4.0	0.003

Table 4: Recombination Line Parameters and Derived Quantities for or NGC 3576

	V (km s ⁻¹)	ΔV (km s ⁻¹)	T_l/T_c	T_e^* (K)
H90 α	-20.0 $\pm$ 0.1	28.7 $\pm$ 0.1	0.105 $\pm$ 0.001	6300 $\pm$ 700
He90 α	-19.3 $\pm$ 0.4	24.8 $\pm$ 0.8	0.011 $\pm$ 0.001	
C90 α	-21.5 $\pm$ 0.9	8.75 $\pm$ 1.8	0.003 $\pm$ 0.001	
H113 β	-22.3 $\pm$ 0.2	33.6 $\pm$ 0.4	0.022 $\pm$ 0.001	7100 $\pm$ 800
H90 α				
<i>Irs1/Irs2</i>	-24.0 $\pm$ 0.1	29.1 $\pm$ 0.1	0.100 $\pm$ 0.001	6500 $\pm$ 700
<i>Irs3</i>	-17.2 $\pm$ 0.1	26.5 $\pm$ 0.1	0.112 $\pm$ 0.001	6400 $\pm$ 700
<i>Irs4</i>	-14.5 $\pm$ 0.1	32.0 $\pm$ 0.3	0.081 $\pm$ 0.001	7200 $\pm$ 800
<i>Irs5</i>	-18.8 $\pm$ 0.2	36.2 $\pm$ 0.4	0.098 $\pm$ 0.002	5500 $\pm$ 600

FIGURE CAPTIONS

Fig. 1 The 3.4 cm continuum image of NGC 3576 is shown at a resolution of $7''$ (0.09 pc at 3.0 kpc). The first positive and negative continuum contours are at 3σ (54 mJy beam^{-1}). Subsequent positive contours are 1.4, 2, 2.8, 4, 5.6, 8, 11.3, 16, 22.6, 32 and 45.3 times the 3σ level. The peak continuum flux density is 4.1 Jy beam^{-1} . The positions of the $10 \mu\text{m}$ sources (Frogel & Person 1974) are indicated with crosses. Errors in the absolute IR positions are $\pm 4''$ (indicated by the cross size). The inset is an integrated profile made over a region of the source where broad recombination lines are detected. The best fit to the spectrum consists of two Gaussians separated by $\sim 25 \text{ km s}^{-1}$. Epoch is B1950.

Fig. 2 The 3.4 cm continuum image of NGC 3603 with the 13 named sources (A to M in order of increasing RA). The continuum data has a spatial resolution of $7''$ (0.21 pc at $D = 6.1 \pm 0.6$ kpc). The first contour is at 5σ (55 mJy beam^{-1}) with subsequent levels at 1.4, 2, 2.8, 4, 5.6, 8, 11.3, 16, 22.6, 32 and 45.3 times the 5σ level. The peak continuum flux density is $0.47 \text{ Jy beam}^{-1}$. The positions of the $10 \mu\text{m}$ sources (Frogel et al. 1977) are indicated with crosses. Errors in the absolute IR positions from this later study are $\pm 10''$ (indicated by the cross size). Epoch is B1950.

Fig. 3 (a) The continuum weighted, integrated line profile for NGC 3576. The recombination lines shown (left to right) are $\text{H}113\beta$, $\text{C}90\alpha$, $\text{He}90\alpha$, and $\text{H}90\alpha$. The solid line is the Gaussian fit, the crosses are the data points, and the dashed line is the residual. The spectral resolution is 2.6 km s^{-1} . (b) The continuum weighted, integrated line profile for the bright southern region in NGC 3603. The recombination lines shown (left to right) are $\text{H}113\beta$, $\text{C}90\alpha$, $\text{He}90\alpha$, and $\text{H}90\alpha$. The solid line is the Gaussian fit, the crosses are the data points, and the dashed line is the residual. The spectral resolution is 2.6 km s^{-1} .

Fig. 4 The continuum weighted, integrated line profiles for the Irs sources in NGC 3576 (Frogel & Persson 1974), indicated in Figure 1. The solid line is the Gaussian fit, the crosses are the data points, and the dashed line is the residual. The spectral resolution is 2.6 km s^{-1} . Figures show (a) Irs 1/2, (b) Irs 3, (c) Irs 4, (d) Irs 5.

Fig. 5 Line parameters of the $\text{H}90\alpha$ line in NGC 3576 overlaid on the 3.4 cm continuum contours. Fits were made to each pixel above the 5σ level in the continuum. (a) The velocity (V_{LSR}) of the $\text{H}90\alpha$ line in NGC 3576 for each pixel is presented in grey-scale with the NGC 3576 3.4 cm continuum contours overlaid. The contour levels are as indicated in Figure 1. The grey scale velocity range is from -35 km s^{-1} to 10 km s^{-1} . (b) The full width at half maximum (ΔV_{FWHM}) of the $\text{H}90\alpha$ line in NGC 3576 for each pixel is presented in false color. The line width range is from 20 km s^{-1} to 50 km s^{-1} . (c) The electron temperature calculated for each pixel in NGC 3576 above 10σ in the continuum image. The temperature (grey scale) ranges from 5000 to 12000 K. The contours start at 4000 K and rise linearly to 12000 K in increments of 1000 K.

Fig. 6 The integrated electron temperature in NGC 3576 for 18 concentric semicircles (width of 4 pixels) centered on the peak in the 3.4 cm continuum image. Temperatures range from 6000 K to 9000 K across the source.


Fig. 7 The integrated line profiles from the 13 individual HII regions in NGC 3603, as indicated in Figure 2. The recombination lines shown (left to right where detected) are H113 β , C90 α , He90 α , and H90 α . The plot shows the Gaussian fit (solid line), data (crosses), and the residual (dashed line). The spectral resolution is 2.6 km s $^{-1}$. Figures 7a to 7m show the integrated profiles of sources NGC 3603 A-M respectively.


Fig. 8 The 3.4 cm continuum contours for NGC 3576. The stellar positions of cluster member stars with infrared excess (Persi et al. 1994) are plotted as crosses. Also plotted are the positions of the star HD 97499 (diamond), and the H $_2$ O (triangle; Caswell 1989) and CH $_3$ OH (box; Caswell et al. 1995) maser positions. Continuum contours are as indicated in Figure 1. Epoch is J2000.


Fig. 9 The 3.4 cm continuum contours in the vicinity of the known stellar cluster HD 97950. The positions of ~ 50 cluster members with measured U , B , and V magnitudes (Melnick, Tapia & Terlevich 1989) are plotted as crosses. Coordinates of the 50 stars were derived from pixel offsets from the known position of Sher 25 (a B1.5Iab supergiant) as given by Brandner et al (1997). Also indicated is the position of HD 97950 (diamond), and H $_2$ O (triangles; Caswell 1989) and OH (box; Caswell 1998) maser positions. Continuum contours are as indicated in Figure 2. Epoch is J2000.

Table 5: Recombination Line Parameters and Derived Quantities for NGC 3603


Source		V (km s ⁻¹)	ΔV (km s ⁻¹)	T_i/T_c	T_e^* (K)	T_e (K)	Y^+	β/α	V_{turb}
Total	H90 α	9.14 $\pm$ 0.1	36.9 $\pm$ 0.25	0.075 $\pm$ 0.001	6600 $\pm$ 500				24 $\pm$ 2
	He90 α	8.9 $\pm$ 1.0	38.9 $\pm$ 2.6	0.009 $\pm$ 0.001			0.13 $\pm$ 0.01		
	C90 α	6.2 $\pm$ 1.9	10.0 (fixed)	0.002 $\pm$ 0.001					
	H113 β	5.9 $\pm$ 0.3	42.6 $\pm$ 0.8	0.026 $\pm$ 0.001	4800 $\pm$ 400			0.39 $\pm$ 0.01	
A	H90 α	8.08 $\pm$ 0.2	26.3 $\pm$ 0.5	0.107 $\pm$ 0.003	6300 $\pm$ 1100	6700 $\pm$ 1200			14 $\pm$ 3
	He90 α	4.0 $\pm$ 1.3	25.0 $\pm$ 3.1	0.019 $\pm$ 0.003			0.17 $\pm$ 0.03		
	H113 β	2.5 $\pm$ 0.8	31.6 $\pm$ 1.9	0.033 $\pm$ 0.003	4900 $\pm$ 900	5200 $\pm$ 1000		0.31 $\pm$ 0.03	
B	H90 α	-3.65 $\pm$ 0.3	28.3 $\pm$ 0.7	0.103 $\pm$ 0.004	6300 $\pm$ 1600	6400 $\pm$ 1600			17 $\pm$ 4
	He90 α	0.9 $\pm$ 2.0	28.8 $\pm$ 4.8	0.016 $\pm$ 0.004			0.16 $\pm$ 0.04		
	C90 α	-12.5 $\pm$ 1.3	10 (fixed)	0.015 $\pm$ 0.003					
	H113 β	-7.4 $\pm$ 1.1	29.0 $\pm$ 2.6	0.029 $\pm$ 0.004	6000 $\pm$ 1500	6100 $\pm$ 1500		0.29 $\pm$ 0.04	
C	H90 α	3.8 $\pm$ 0.7	33.2 $\pm$ 1.7	0.104 $\pm$ 0.007	4300 $\pm$ 1700	4200 $\pm$ 1700			22 $\pm$ 9
	He90 α	9.1 $\pm$ 4.3	44.1 $\pm$ 10	0.020 $\pm$ 0.006			0.26 $\pm$ 0.10		
	H113 β	-5.2 $\pm$ 4.1	49.1 $\pm$ 10	0.023 $\pm$ 0.006	4400 $\pm$ 1700	4300 $\pm$ 1700		0.32 $\pm$ 0.11	
D	H90 α	-0.80 $\pm$ 0.4	35.9 $\pm$ 1.1	0.078 $\pm$ 0.003	6400 $\pm$ 2000	6800 $\pm$ 2100			23 $\pm$ 7
	He90 α	1.1 $\pm$ 2.2	32.8 $\pm$ 5.6	0.014 $\pm$ 0.003			0.16 $\pm$ 0.05		
	H113 β	-1.7 $\pm$ 1.0	31.4 $\pm$ 2.6	0.029 $\pm$ 0.003	5700 $\pm$ 1800	6100 $\pm$ 1900		0.33 $\pm$ 0.05	
E	H90 α	1.9 $\pm$ 0.2	41.9 $\pm$ 0.5	0.066 $\pm$ 0.001	6700 $\pm$ 600	8500 $\pm$ 800			27 $\pm$ 3
	He90 α	-2.2 $\pm$ 1.5	36.9 (fixed)	0.008 $\pm$ 0.001			0.11 $\pm$ 0.01		
	H113 β	-2.3 $\pm$ 0.6	46.5 $\pm$ 1.4	0.022 $\pm$ 0.001	5300 $\pm$ 500	6700 $\pm$ 600		0.36 $\pm$ 0.02	
F	H90 α	11.5 $\pm$ 0.8	52.7 $\pm$ 2.2	0.056 $\pm$ 0.003	6500 $\pm$ 3300	7000 $\pm$ 3600			36 $\pm$ 19
	He90 α	27.6 $\pm$ 2.7	22.7 $\pm$ 6.9	0.011 $\pm$ 0.005			0.08 $\pm$ 0.04		
	H113 β	5.9 $\pm$ 2.4	59.9 $\pm$ 6.8	0.020 $\pm$ 0.003	4700 $\pm$ 2400	5100 $\pm$ 2600		0.40 $\pm$ 0.08	
G	H90 α	15.5 $\pm$ 0.4	25.6 $\pm$ 1.0	0.115 $\pm$ 0.006	6200 $\pm$ 2200	7900 $\pm$ 2800			13 $\pm$ 5
	He90 α	20.0 $\pm$ 1.6	17.2 $\pm$ 3.8	0.024 $\pm$ 0.001			0.14 $\pm$ 0.05		
	H113 β	8.3 $\pm$ 1.2	31.2 $\pm$ 3.1	0.040 $\pm$ 0.006	4400 $\pm$ 1500	5600 $\pm$ 1900		0.43 $\pm$ 0.07	
H	H90 α	11.8 $\pm$ 0.1	32.2 $\pm$ 0.3	0.082 $\pm$ 0.001	6900 $\pm$ 1200	8300 $\pm$ 1400			19 $\pm$ 3
	He90 α	12.4 $\pm$ 0.9	29.3 $\pm$ 2.2	0.011 $\pm$ 0.001			0.12 $\pm$ 0.02		
	C90 α	8.2 $\pm$ 1.6	10.0 (fixed)	0.003 $\pm$ 0.001					
	H113 β	8.5 $\pm$ 0.4	37.9 $\pm$ 0.9	0.028 $\pm$ 0.001	5100 $\pm$ 900	6300 $\pm$ 1100		0.40 $\pm$ 0.02	
I	H90 α	15.9 $\pm$ 0.2	34.8 $\pm$ 0.6	0.068 $\pm$ 0.002	7900 $\pm$ 2200	9300 $\pm$ 2600			21 $\pm$ 6
	He90 α	7.9 $\pm$ 3.1	44.9 $\pm$ 8.0	0.006 $\pm$ 0.001			0.11 $\pm$ 0.03		
	H113 β	13.4 $\pm$ 0.7	39.6 $\pm$ 1.8	0.023 $\pm$ 0.002	5900 $\pm$ 1600	6800 $\pm$ 1800		0.39 $\pm$ 0.03	
J	H90 α	7.3 $\pm$ 0.3	39.9 $\pm$ 0.9	0.085 $\pm$ 0.003	5700 $\pm$ 1600	5800 $\pm$ 1600			26 $\pm$ 7
	He90 α	2.9 $\pm$ 3.6	59.0 $\pm$ 9.7	0.011 $\pm$ 0.003			0.18 $\pm$ 0.05		
	H113 β	5.3 $\pm$ 1.2	47.8 $\pm$ 2.9	0.031 $\pm$ 0.003	3600 $\pm$ 1000	3700 $\pm$ 1000		0.43 $\pm$ 0.05	
K	H90 α	5.5 $\pm$ 0.3	40.8 $\pm$ 0.7	0.093 $\pm$ 0.002	5100 $\pm$ 1000	5000 $\pm$ 1000			28 $\pm$ 6
	He90 α	7.9 $\pm$ 1.9	31.2 $\pm$ 4.7	0.012 $\pm$ 0.003			0.10 $\pm$ 0.02		
	H113 β	3.8 $\pm$ 0.7	43.8 $\pm$ 1.9	0.037 $\pm$ 0.002	3600 $\pm$ 700	3500 $\pm$ 700		0.43 $\pm$ 0.03	
L	H90 α	6.4 $\pm$ 0.6	38.2 $\pm$ 1.5	0.142 $\pm$ 0.008	3800 $\pm$ 1000	3700 $\pm$ 1000			26 $\pm$ 7
	He90 α	5.7 $\pm$ 4.5	35.0 (fixed)	0.018 $\pm$ 0.004			0.11 $\pm$ 0.03		
	H113 β	2.7 $\pm$ 1.3	32.9 $\pm$ 3.1	0.063 $\pm$ 0.008	2900 $\pm$ 800	2800 $\pm$ 800		0.37 $\pm$ 0.07	
M	H90 α	4.6 $\pm$ 0.5	37.3 $\pm$ 2.1	0.115 $\pm$ 0.009	3600 $\pm$ 1300	3500 $\pm$ 1300			26 $\pm$ 10
	He90 α	8.5 $\pm$ 6.3	33.8 (fixed)	0.015 $\pm$ 0.005			0.11 $\pm$ 0.04		
	H113 β	4.3 $\pm$ 2.1	44.9 $\pm$ 5.4	0.051 $\pm$ 0.009	2600 $\pm$ 1000	2500 $\pm$ 1000		0.53 $\pm$ 0.12	













Velocity (km/s)







Velocity (km/s)













$T(L)/T(C)$



Velocity (km/s)

